



Bayer CropScience LP  
P.O. Box 12014  
2 T.W. Alexander Drive  
Research Triangle Park, North Carolina 27709  
1-866-99BAYER (1-866-992-2937)

## Velum® Prime

In-furrow at-plant applications in potato for the suppression of nematodes, white mold, and early blight.

For use only on potatoes to be consumed in the US, Canada, and Mexico. All potato by-products must be used in the US, Canada, and Mexico.

EPA Reg. No. 264-1078

## FIFRA 2(ee) Recommendation

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE(S) OF ALABAMA, ARKANSAS, COLORADO, FLORIDA, GEORGIA, IDAHO, INDIANA, KANSAS, LOUISIANA, MAINE, MICHIGAN, MINNESOTA, MISSISSIPPI, MONTANA, NORTH CAROLINA, NORTH DAKOTA, NEBRASKA, NEW JERSEY, OREGON, PENNSYLVANIA, SOUTH CAROLINA, TENNESSEE, UTAH, VIRGINIA WASHINGTON, WISCONSIN**

**FIFRA SECTION 2(EF) RECOMMENDATION:** This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label.

Potato		
Pest Suppression	Application Rate	Product Instructions
Nematodes  Early blight ( <i>Alternaria solani</i> )  White mold ( <i>Sclerotinia sclerotiorum</i> )	6.5 fl oz/acre	<b>Soil Applications:</b> Apply specified dosage as an in-furrow spray during planting directed on or below seed. Despite suppression of root-knot nematode, tuber quality may not be adequately protected. If root-knot nematode is a severe economic pest, the use of other suppression measures should also be employed
<b>Restrictions:</b> <ul style="list-style-type: none"><li>Do not apply more than 13.7 fl oz of VELUM PRIME (0.446 lbs Fluopyram) per acre per year, regardless of formulation or method of application (soil or foliar).</li><li>Do not apply VELUM PRIME within 7 days of harvest.</li><li>To limit the potential for development of disease resistance to this fungicide class, do not make more than 2 sequential applications of VELUM PRIME or any Group 7-containing fungicide before rotating with a fungicide from a different Group.</li><li>Tops or greens from treated product can be utilized for food or feed.</li><li>The grazing of livestock in treated areas within 7 days of application is prohibited.</li></ul>		

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.  
For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).

*As with any crop-protection product, always read and follow label directions.  
For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*